

# RELATIONS

## CIS002-2 COMPUTATIONAL ALGEBRA AND NUMBER THEORY

David Goodwin

david.goodwin@perisic.com



11:00, Tuesday 29<sup>th</sup> November 2011

# OUTLINE

## ① RELATIONS

Relation

Reflexive relation

Symmetric relation

Antisymmetric relation

Transitive relation

Partial order

Inverse

Composition

## ② EQUIVALENCE RELATIONS

Equivalence relation

Equivalence classes

## ③ CLASS EXERCISES

# OUTLINE

## ① RELATIONS

Relation

Reflexive relation

Symmetric relation

Antisymmetric relation

Transitive relation

Partial order

Inverse

Composition

## ② EQUIVALENCE RELATIONS

Equivalence relation

Equivalence classes

## ③ CLASS EXERCISES

# RELATIONS

A (binary) **relation**  $R$  from a set  $X$  to a set  $Y$  is a subset of the Cartesian product  $X \times Y$ . If  $(x, y) \in R$ , we write  $xRy$  and say that  $x$  is related to  $y$ . If  $X = Y$ , we call  $R$  a (binary) relation on  $X$ .

A function is a special type of relation. A function  $f$  from  $X$  to  $Y$  is a relation from  $X$  to  $Y$  having the properties:

- The domain of  $f$  is equal to  $X$ .
- For each  $x \in X$ , there is exactly one  $y \in Y$  such that  $(x, y) \in f$

## RELATIONS - EXAMPLE

Let

$$X = \{2, 3, 4\} \quad \text{and} \quad Y = \{3, 4, 5, 6, 7\}$$

If we define a relation  $R$  from  $X$  to  $Y$  by

$$(x, y) \in R \quad \text{if } x \mid y$$

we obtain

$$R = \{(2, 4), (2, 6), (3, 3), (3, 6), (4, 4)\}$$

# RELATIONS - REFLEXIVE

A relation  $R$  on a set  $X$  is **reflexive** if  $(x, x) \in R$ , if  $(x, y) \in R$  for all  $x \in X$ .

# RELATIONS - SYMMETRY

A relation  $R$  on a set  $X$  is **symmetric** if for all  $x, y \in X$ , if  $(x, y) \in R$  then  $(y, x) \in R$ . In symbols we can write

$$\forall x \forall y [(x, y) \in R] \rightarrow [(y, x) \in R]$$

# RELATIONS - ANTISYMMETRY

A relation  $R$  on a set  $X$  is **antisymmetric** if for all  $x, y \in X$ , if  $(x, y) \in R$  and  $(y, x) \in R$ , then  $x = y$ . In symbols we can write

$$\forall x \forall y [(x, y) \in R \wedge (y, x) \in R] \rightarrow [x = y]$$



# RELATIONS - TRANSITIVE

A relation  $R$  on a set  $X$  is **transitive** if for all  $x, y, z \in X$ , if  $(x, y) \in R$  and  $(y, z) \in R$ , then  $(x, z) \in R$ . In symbols we can write

$$\forall x \forall y \forall z [(x, y) \in R \wedge (y, z) \in R] \rightarrow [(x, z) \in R]$$

# RELATIONS - PARTIAL ORDER

A relation  $R$  on a set  $X$  is a **partial order** if  $R$  is reflexive, antisymmetric, and transitive. If  $R$  is a partial order on a set  $X$ , the notation  $x \preceq y$  is sometimes used to indicate that  $(x, y) \in R$ .

Suppose that  $R$  is a partial order on a set  $X$ . If  $x, y \in X$  and either  $x \preceq y$  or  $y \preceq x$  we say  $x$  and  $y$  are **comparable**. If  $x, y \in X$  and  $x \not\preceq y$  and  $y \not\preceq x$  we say  $x$  and  $y$  are **incomparable**. If every pair of elements in  $X$  is comparable, we call  $R$  a **total order**.

# RELATIONS - INVERSE

A  $R$  be a relation from  $X$  to  $Y$ . The **inverse** of  $R$ , denoted  $R^{-1}$ , is the relation from  $Y$  to  $X$  defined by

$$R^{-1} = \{(y, x) \mid (x, y) \in R\}$$

# RELATIONS - COMPOSITION

Let  $R_1$  be a relation from  $X$  to  $Y$  and  $R_2$  be a relation from  $Y$  to  $Z$ . The **composition** of  $R_1$  and  $R_2$ , denoted  $R_2 \circ R_1$ , is the relation from  $X$  to  $Z$  defined by

$$R_2 \circ R_1 = \{(x, z) \mid (x, y) \in R_1 \text{ and } (y, z) \in R_2 \text{ for some } y \in Y\}$$

# OUTLINE

## ① RELATIONS

Relation

Reflexive relation

Symmetric relation

Antisymmetric relation

Transitive relation

Partial order

Inverse

Composition

## ② EQUIVALENCE RELATIONS

Equivalence relation

Equivalence classes

## ③ CLASS EXERCISES

**THEOREM**

*Let  $S$  be a partition of the set  $X$ . Define  $xRy$  to mean that for some set  $S$  in  $S$ , both  $x$  and  $y$  belong to  $S$ . Then  $R$  is reflexive, antisymmetric, and transitive.*

---

A partition of a set  $X$  is a collection  $S$  of nonempty subsets of  $X$  such that every element in  $X$  belongs to exactly one member of  $S$ .

# EQUIVALENCE RELATION

A relation that is reflexive, symmetric, and transitive on a set  $X$  is called an **equivalence relation** on  $X$ .

## THEOREM

Let  $R$  be an equivalence relation on a set  $X$ . For each  $a \in X$ , let

$$[a] = \{x \in X \mid xRa\}$$

(In words,  $[a]$  is a set of all elements in  $X$  that are related to  $a$ .)

Then

$$\mathcal{S} = \{[a] \mid a \in X\}$$

is a partition of  $X$ .



# EQUIVALENCE CLASSES

Let  $R$  be an equivalence relation on a set  $X$ . The sets  $[a]$  are called the **equivalence classes** of  $X$  given by the relation  $R$ .

## THEOREM

*Let  $R$  be an equivalence relation on a finite set  $X$ . If each equivalence class has  $r$  elements, there are  $|X|/r$  equivalence classes.*

# OUTLINE

## ① RELATIONS

- Relation
- Reflexive relation
- Symmetric relation
- Antisymmetric relation
- Transitive relation
- Partial order

- Inverse
- Composition

## ② EQUIVALENCE RELATIONS

- Equivalence relation
- Equivalence classes

## ③ CLASS EXERCISES

## EXERCISES

- 1 Determine whether the following relation defined on the set of positive integers is reflexive, symmetric, antisymmetric, transitive, and / or a partial order
  - A  $(x, y) \in R$  if  $2 \mid x + y$
  - B  $(x, y) \in R$  if  $3 \mid x + y$
- 2 Give an example of a relation on  $\{1, 2, 3, 4\}$  that is reflexive, not antisymmetric, and not transitive.

## EXERCISES

- 3 Suppose that  $R$  is a relation on  $X$  that is symmetric and transitive but not reflexive. Suppose also that  $|X| \geq 2$ . Define the relation  $\bar{R}$  on  $X$  by

$$\bar{R} = X \times X - R$$

Which of the following must be true? For each false statement, provide a counterexample

- A  $\bar{R}$  is reflexive
- B  $\bar{R}$  is symmetric
- C  $\bar{R}$  is not antisymmetric
- D  $\bar{R}$  is transitive

## EXERCISES

- 4 Is the relation

$$\{(1, 1), (1, 2), (2, 2), (4, 4), (2, 1), (3, 3)\}$$

an equivalence relation on  $\{1, 2, 3, 4\}$ ? Explain.

- 5 Given the relation

$$\{(1, 1), (2, 2), (3, 3), (4, 4), (1, 2), (2, 1), (3, 4), (4, 3)\}$$

is an equivalence relation on  $\{1, 2, 3, 4\}$ , find  $[3]$ , the equivalence class containing 3. How many distinct equivalence classes are there?

## EXERCISES

- 6 Find the equivalence relation (as a set of ordered pairs) on  $\{a, b, c, d, e\}$ , whose equivalence classes are  $\{a\}$ ,  $\{b, d, e\}$ ,  $\{c\}$ .
- 7 Let  $R$  be the relation defined on the set of eight-bit strings by  $s_1 R s_2$  provided that  $s_1$  and  $s_2$  have the same number of zeros.
- A Show that  $R$  is an equivalence relation.
- B How many equivalence classes are there?
- C List one member of each equivalence class.