

Betti numbers and designs of experiments

Hugo Maruri-A., Eduardo Sáenz, Henry Wynn

School of Mathematical Sciences
Queen Mary, University of London



WOGAS 3, Warwick University, 7th April 2011

Betti numbers and designs of experiments

Algebraic techniques in experimental designs were pioneered by Pistone and Wynn (1996). Recently, the models identified with those techniques have been observed to be of a low average degree. In other words, the hierarchical monomials forming the model are those that, while still being linearly independent (moduli design ideal), are of low weighted degree. An interesting characterization of the complexity of such models is given by the number of generators of the leading term ideal, i.e. the generator set of those monomials not in the model. This number is computed by the Betti numbers associated to the ideal of leading terms.

I will present some examples related to generic designs which highlight that, most of the models are simultaneously of minimal aberration and have maximal Betti numbers. Some further extensions to designs which are fractions of factorial designs will be discussed.

1. The algebraic approach to identifiability

The algebraic approach

Identifications of models from a given design can be achieved by computational commutative algebra techniques (Pistone and Wynn, 1996), with the design \mathcal{D} considered as an algebraic variety (i.e. a solution of a system of polynomial equations).

- Models are identified
- Confounding relations between factors are generalized

This approach has been worked in the context of industrial experimentation (Halliday et al., 1996), mixture designs (Maruri et al., 2006).

The collection of algebraic models (Caboara et al., 1997) has been identified to be of low average degree (Bernstein et al., 2010). This is a characterisation of the centroid of the model. Recent work characterises instead the border of the model (Maruri et al., 2010).

Rings, polynomial division (Cox et al., 1996)

- $\mathbb{R}[x] = \mathbb{R}[x_1, \dots, x_d]$ the polynomial ring.
- The ideal generated by a finite set of points $\mathcal{D} \subset \mathbb{R}^d$ is $I(\mathcal{D}) = \{f \in \mathbb{R}[x] : f(x) = 0, x \in \mathcal{D}\} \subset \mathbb{R}[x]$.
- A term order τ is a total ordering in monomials in $T^d = \{x^\alpha : \alpha \in \mathbb{Z}_{\geq 0}^d\}$, compatible with monomial simplification: i) $x^\alpha \succ 1, \alpha \neq \mathbf{0}$, ii) $x^\alpha \succ x^\beta \Rightarrow x^{\alpha+\gamma} \succ x^{\beta+\gamma}$ for $x^\alpha, x^\beta, x^\gamma \in T^d$.
- A Gröbner basis G_τ is a finite subset of $I(\mathcal{D})$ such that $\langle \text{LT}(g) : g \in G_\tau \rangle = \langle \text{LT}(f) : f \in I(\mathcal{D}) \rangle$.
- For any $f \in \mathbb{R}[x]$, unique remainder r in division of f by $I(\mathcal{D})$


$$f = \sum_{g \in G_\tau} gh + r \quad (1)$$

Quotient rings (Cox et al., 1996)

- $\mathbb{R}[\mathcal{D}]$ is the collection of polynomial functions $\phi : \mathcal{D} \mapsto \mathbb{R}$.
- The elements of $\mathbb{R}[\mathcal{D}]$ are in one to one correspondence with equivalence classes of polynomials modulo $I(\mathcal{D})$ and we have an isomorphism $\mathbb{R}[\mathcal{D}] \sim \mathbb{R}[x]/I(\mathcal{D})$.
- A basis for $\mathbb{R}[x]/I(\mathcal{D})$ is given by those monomials that cannot be divided by any of $\text{LT}(g)$ for $g \in G_\tau$.
- The remainder in Eq. (1) is known as the normal form of f (modulo $I(\mathcal{D})$), i.e. $\text{NF}(f) = r$.

Generalised confounding (Pistone and Wynn, 1996)

- Design \mathcal{D} , n points, d factors.
- Study the \mathcal{D} through the design ideal $I(\mathcal{D}) \subset \mathbb{R}[x]$.
- The support for a model is given by those monomials not divisible by the leading terms of the RGröbner basis $G_\tau \subset I(\mathcal{D})$.



$$G_\tau = \{\underline{x_1^2} + 2x_1x_2 + x_2^2 - x_1 - x_2, \underline{x_2^3} - x_2, \underline{x_1x_2^2} - x_1x_2 - x_2^2 + x_2\}$$

- Exact polynomial interpolator = saturated regression model.
- Hierarchical polynomial model: staircases.
- Link with aliasing/confounding $f(x) = g(x), x \in \mathcal{D}$.


Examples

- Factorial design 2^d with levels ± 1 . For any term ordering, its design ideal $I(\mathcal{D})$ has Gröbner basis $G_\tau = \{x_i^2 - 1, i = 1, \dots, d\}$ and identifies the model $\{1, x_1\} \times \dots \times \{1, x_d\}$
 - Indicator function blends naturally to create the ideal of a design fraction, e.g. the indicator $(x_1 - x_2)(x_2 - x_3)$ removes the treatments $\pm(1, -1, 1)$ from the 2^3 design. The fraction \mathcal{F} has six runs and for the standard term order in CoCoA, the model identified is $\{1, x_1, x_2, x_3, x_1x_3, x_2x_3\}$.
 - Confounding by normal form: $\text{NF}(x_1x_2x_3) = x_1 - x_2 + x_3$.
 - Technique applicable to any design whose points have continuous factors: LH, RSM, optimal. Adaptable to other structures: block, row-column, ...
- ... linear independence with a term order, but much more!

A column selection algorithm (Babson et al., 2003)

- Compute the design model matrix for the set of terms V_n^d .
- Using a term ordering \succ_w , order the columns of the matrix.
- Pick the first n columns which form a linearly independent set.

$$\begin{array}{ccccccc}
 1 & x_1 & x_2 & x_1^2 & x_1x_2 & x_2^2 & x_1^3 & \cdots \\
 \hline
 1 & 0 & 0 & 0 & 0 & 0 & 0 & \\
 1 & 1 & 0 & 1 & 0 & 0 & 1 & \\
 1 & 0 & 1 & 0 & 0 & 1 & 0 & \\
 1 & 1 & -1 & 1 & -1 & 1 & 1 & \\
 1 & -1 & 1 & 1 & -1 & 1 & -1 &
 \end{array}$$




$$V_n^d := \{x \in \mathbb{Z}_{\geq 0}^d : \prod_{i=1}^d (x_i + 1) \leq n\}$$

By row elimination, the methodology retrieves G_w for $I(\mathcal{D})$.

It is a variation of the FGLM algorithm for change of basis Faugere *et al.* (1993).


2. The fan of a design

- As we scan over all possible term orders, we obtain the algebraic fan of \mathcal{D} , see (Caboara *et al.*, 1997 and Maruri, 2007).
- Not all identifiable hierarchical models belong to the algebraic fan, i.e. $\emptyset \subset A \subseteq S \subseteq \mathcal{C}_{d,n}$.



$$A = \{ \dot{\vdots} \quad , \quad \vdots \dot{\vdots} \}, S \setminus A = \{ \dot{\vdots} \dots \}$$

- The models in A correspond to the vertexes of the state polyhedron $\mathcal{S}(I)$, e.g. we add up the exponent vectors for $L = \{1, x_1, x_2, x_1x_2, x_2^2\}$, $\bar{\alpha}_L = \sum_L \alpha = (2, 4)$.



$$\mathcal{S}(I) = \text{conv}(\bar{\alpha}_L : L \in A) + \mathbb{R}_{\geq 0}^d$$

3. Linear aberration

Linear aberration of model L

- Taking the motivation from the concept of aberration, we want to fill out lower degrees before higher:


$$A(w, L) = \frac{1}{n} \sum w_i \bar{\alpha}_{L_i}$$

$$w_i \geq 0, \sum w_i = 1.$$

Theo. There exists w such that $A(w, L)$ is minimised by an algebraic model.

Proof. Use LP arguments for the lower boundary of $\mathcal{S}(I)$.

- **Generic** designs minimise $A(w, L)$ over $\mathcal{C}_{d,n}$ and all vectors w .
- For generic designs, algebraic models are corner cut models [??]


Corner cut model  $\{1, x_1, x_2, x_1x_2\}$ is not corner cut

Linear aberration and algebraic models

- The state polytope summarises information about linear aberration, i.e. its vertexes correspond to models that minimise $A(w, L)$ over the set of identifiable hierarchical models \mathcal{S} .
- The vertexes of $\mathcal{S}(I)$ correspond to algebraic models A .
- The (minimum) aberration of designs can be compared through their state polytopes.
- However, there may be non-algebraic models on the lower boundary (and thus minimising $A(w, L)$ for some w) or in the interior of $\mathcal{S}(I)$.

Example aberration 1


Central composite design (CCD, Box, 1957) with $d = 2$, $n = 9$ and axial distance = $\sqrt{2}$



Algebraic = $\{1, x_1, x_1^2, x_1^3, x_1^4, x_2, x_1x_2, x_1^2x_2, x_2^2\}$ and its conjugate

Example aberration 2

Consider $\mathcal{D} = \{(0, 0), (1, 1), (2, 2), (3, 4), (5, 7), (11, 13), (\alpha, \beta)\}$, with $(\alpha, \beta) \approx (1.82997, 1.82448)$ (Onn, 1999)



\Rightarrow The set of algebraic models can be larger in size than the set of corner cut models. However, corner cut models are always of lowest possible degree over all vectors $c \neq 0$.

Minimal aberration (Bernstein et al. (2010) [??,??])

L a model support, $w > 0$ vector of weights, $\sum w_i = 1$

Compute the aberration: $A(w, L) = \frac{1}{n} \sum w_i \bar{\alpha}_{L_i}$

Minimal aberration II (Bernstein et al. (2010) [??,??])

For fixed w , define

$$A^* = \min_{L \in \mathcal{L}} A(w, L).$$

This value is achieved for algebraic models in a generic design.


Theorem. In a generic design, minimal aberration A^* obeys the following bounds:

$$A^+ - 1 \leq A^* \leq A^+ + 1$$

with $A^+ = (nd!)^{\frac{1}{d}} \frac{d}{d+1} g(w)$ and $g(w) = \sqrt[d]{w_1 \cdots w_d}$

Proof.

Define equivalent simplex $S(w)$ ($\int S = n$) and lower and upper cells \underline{Q} and \bar{Q} .




Aberration over S is $E(w^T X)$ with $X \sim U(S)$. The bounds follow from the following inequalities:


$$A(w, \underline{S}(w)) \leq A(w, \underline{Q}) \leq A(w, S(w)) \leq A(w, \overline{Q}),$$

and noting that $A(w, \underline{S}(w)) = A(w, S(w)) - 1$. We denote A^+ for $A(w, S(w))$.


Example: minimal aberration $d = 2$



$$n = 4$$



$$n = 20$$



$$n = 100$$

The graphs include A^* , A^+ and $A^+ \pm 1$ of Theorem; also approximate \tilde{A} is shown.

4. The border of the model

For a polynomial ideal I , the *ideal of leading terms* $LT(I)$ is the monomial ideal generated by the leading terms of polynomials in I . In what remains of the talk, I denotes a monomial ideal.

The Hilbert function $H_{R/I}$ counts the number of monomials not in I , for each degree. E.g. for $I = \langle x^3, y^2 \rangle$ we have $H_{R/I} = 1, 2, 2, 1$.

The generating function for those terms is the *multigraded Hilbert series* $HS_{R/I}$. In the current example $HS_{R/I} = 1 + x + y + x^2 + xy + x^2y$.

We can however compute the Hilbert function and the Hilbert series for terms in I , and we have the following equality for HS :

$$\sum_{\alpha \geq 0} x^\alpha = \sum_{\alpha \in I} x^\alpha + \sum_{\alpha \notin I} x^\alpha$$
$$\frac{1}{\prod_{i=1}^d (1 - x_i)} = HS_I + HS_{R/I}$$

The table of Betti numbers describes the composition of (numerators of) sums, i.e. entry (i, j) contains number of terms of degree $i + j$.

$$\frac{1}{(1-x)(1-y)} = \frac{y^2+x^3-x^3y^2}{(1-x)(1-y)} \cup (1+x+y+x^2+xy+x^2y)$$

	0	1	2
0:	1	-	-
1:	-	1	-
2:	-	1	-
3:	-	-	1
Tot:	2	1	1

$$\frac{1}{(1-s)^2} = \frac{s^2+s^3-s^5}{(1-s)^2} + (1+2s+2s^2+s^3)$$

	0	1	2
0:	1	-	-
1:	-	1	-
2:	-	1	-
3:	-	-	1
Tot:	1	2	1

CoCoA code for computing Hilbert function H , Hilbert series HS and Betti table.

```
Use T:=Q[x,y];  
I:=Ideal(x^3,y^2);  
I;  
Hilbert(T/I);  
HilbertSeries(T/I);  
BettiDiagram(I);  
BettiDiagram(T/I);
```

		0	1
...	2:	1	-
...	3:	1	-
	4:	-	1
	Tot:	2	1

		0	1
..	2:	1	-
.....	3:	1	1
	4:	1	1
	Tot:	3	2

		0	1
.	2:	1	-
.....	3:	1	1
	4:	1	1
	Tot:	3	2

		0	1
.	2:	-	-
.....	3:	4	3
	Tot:	4	3

We can only compare tables of Betti numbers for ideals that have the same Hilbert function. In such case, the following theorem guarantees existence of an ideal (called *lex segment ideal*) that attains maximal Betti numbers.

Theorem[Bigatti-Hulett] Let $I \subseteq R$ and L be the lex ideal such that $H_{R/I} = H_{R/L}$. Then $\beta_{i,j}(R/L) \geq \beta_{i,j}(R/I)$ for all i, j .


We want to describe the (borders of) models in the algebraic fan of a design, and we first study generic designs. For generic designs, the models in the algebraic fan are corner cuts.

In two dimensions, the relation between ideals generated by corner cuts staircases and lex-segment ideals is one-to-one. In other words, the models in the algebraic fan are precisely those that maximise Betti numbers (Maruri *et al.*, 2011).


For more than two dimensions, the relationship between lex-segment ideals and ideals which are the complement of corner cut staircases and

is not necessarily one-to-one. For some cases, the ideal of a corner cut model may attain maximal Betti number despite not being a lex-segment ideal, while in other cases it may not attain maximal Betti numbers.

Example. Generic design $n = 7, d = 3$. Fan with 36 models, of which 3 are not lex segment, yet they still attain maximal Betti numbers.




State polytope



Fan (dual of SP)

We give some (necessary) conditions for identification of corner cut models whose ideals are lex segment ideals. The construction builds a trajectory $w = (C - \gamma^{d-1}, C - \gamma^{d-2}, \dots, C - 1)$ in the dual of the state polytope. The initial and final points of the trajectory are lex-segment ideals.



Trajectory ($\gamma = 4$) and the normal fan of the corner cut polytope,
 $d = 3, n = 12$.

5. Betti numbers, the squarefree case

Design whose points are fractions of factorial design with two levels 2^d have important role in experimentation.

Fractions can be selected to satisfy orthogonality conditions and thus not only economy of runs is achieved, but also independent estimation.

Models are (hierarchical) squarefree models and thus they can be seen as simplicial complexes.

For a simplicial complex (model) Δ , we construct the *Stanley Reisner* ideal I_Δ , which will be used to analyze the complexity of the border of Δ . Those ideals will be compared against (square free) lex segment ideals.

5. Betti numbers: An example with Plackett-Burman designs

- Small fractions of 2^d with d factors and $n = d + 1$ runs.
- Designs constructed by circular shifts of a generator, available for $d = 7, 11, 15, 19, 23, \dots$
- PB designs possess a complicated aliasing table, but they have an orthogonal design-model matrix for the linear model with all factors

$$E(y) = \beta_0 + \sum_{i=1}^d \beta_i x_i \quad (2)$$

- PB designs are a popular choice for screening in a first stage of experimentation.

We study their algebraic fan and describe the structure of models with the aid of Betti numbers.

PB8

Consider a Plackett-Burman (PB8) design with eight runs, seven factors a, b, c, d, e, f, g and generator $+ - - + - + +$.

	a	b	c	d	e	f	g
1	1	-1	-1	1	-1	1	1
2	1	1	-1	-1	1	-1	1
3	1	1	1	-1	-1	1	-1
4	-1	1	1	1	-1	-1	1
5	1	-1	1	1	1	-1	-1
6	-1	1	-1	1	1	1	-1
7	-1	-1	1	-1	1	1	1
8	-1	-1	-1	-1	-1	-1	-1

PB8 (cont.)

The design PB8 has 218 models in its fan, which are summarized in Table 2, where representatives of six equivalence classes (up to permutation of factors) are shown.







 $1 + 5s + 2s^2$ (84)	\prec_2 (DegRevLex)  $1 + 7s$ (1)	\prec_3 (Lex)  $1 + 3s + 3s^2 + s^3$ (28)
 $1 + 6s + s^2$ (21)	\prec_1 (Block)  (56) $1 + 4s + 3s^2$ (28)	

Table 2: Equivalence classes of models Δ and corresponding Hilbert Series for PB8.

PB8: Comparing models

Betti table									Model
	0	1	2	3	4	5	6	7	
0:	1	3	3	1	-	-	-	-	
1:	-	7	29	48	40	17	3	-	
2:	-	-	6	26	45	39	17	3	
Tot:	1	10	38	75	85	56	20	3	

	0	1	2	3	4	5	6	7	
0:	1	3	3	1	-	-	-	-	
1:	-	7	30	52	47	24	7	1	
2:	-	1	10	33	52	43	18	3	
Tot:	1	11	43	86	99	67	25	4	

Disconnected model \prec_2 DegRevLex:							
	0	1	2	3	4	5	6
0:	1	-	-	-	-	-	-
1:	-	21	70	105	84	35	6
Tot:	1	21	70	105	84	35	6


PB12

This design has 12 runs in 11 factors and generator
+ + - + + + - - - + -.

The algebraic fan of PB12 is very complex, showing a rich variety of simplicial models.

Despite its enormous size (around 3×10^5), models have been classified in nineteen classes (up to permutations of variables), which in turn share only ten distinct Hilbert Series.

PB12 (cont.)



6. Final comments

Betti numbers provide a description of the model border. Maximal Betti numbers are related to models of low degree (low aberration). However, differently to aberration, we can only compare models (ideals) that have the same Hilbert function.

For generic designs, we found that corner cut models fall into three cases:

- a) when they are lex segment and thus they have maximal Betti numbers,
- b) they are not lex segment yet they still have maximal Betti,
- c) they are not lex segment nor with maximal Betti.

For fractions of 2^d , we have only found classes a) and c). Work in progress...

7. References

1. Babson, Onn, Thomas (2003). *Adv. Appl. Math.* **30**(3), 529-544.
2. Bernstein *et al.* (2007). *Minimal aberration* arXiv:0808.3055.
3. Caboara *et al.* (1997). Res. Rep. 33, Univ. of Warwick.
4. Cheng, Mukerjee (1998). *Ann. Statist.* **26**(6), 2289-2300.
5. Cox, Little, O'Shea (1996). *Ideals, varieties and algorithms*.
6. Faugere *et al.* (1993). *J. Symb. Comput.* **16**, 329-344.
7. Gritzmann, Sturmfels (1993). *SIAM JDM* **6**(2), 246-269.
8. Jensen (2005). Gfan, software.
9. Lee *et al.* (2008) *SIAM JDM* **22**(3), 901-919.
10. Maruri (2006). *RIMUT* **37**(1-2), 95-119.
11. Maruri (2007). Ph.D. Thesis, Univ. of Warwick.
12. Maruri, Sáenz, Wynn (2011). Submitted to ANAI.
13. Maruri, Notari, Riccomagno (2007). *Stat. Sin.* **17**, 1417-1440.

14. Onn, Sturmfels (1999). *Adv. Appl. Math.* **23**, 29-48.
15. Pistone and Wynn (1996). *Biometrika* **83**(3), 653-666.
16. Pistone *et al.* (2000). *Algebraic Statistics*. CRC.
17. Plackett, Burman (1946). *Biometrika* **33**, 305-325.

Thank you!