



## News from the $W$ boson

Jan Kretzschmar, University of Liverpool  
Seminar at Warwick University

18.1.2024


# 40 years since the discovery of the $W$ and $Z$ bosons



- ▶ Force carriers of the electro-weak interaction, acquire their mass from the interaction with the Higgs boson (that was only discovered  $\sim 30$  years later)
- ▶ Discovered in  $p\bar{p}$  collisions at CERN SPS

# W and Z Physics

- ▶ Electroweak sector (almost) completely fixed with just three parameters, e.g.  $\alpha$ ,  $G_F$ ,  $m_Z$
- ▶ Dedicated  $e^+e^- \rightarrow Z$  program at LEP (and SLAC)  $\sim 20$  million Z bosons, but only few W bosons





- ▶ W-boson mass related to other SM parameters

$$m_W^2 \sin^2 \theta_W = \frac{\pi \alpha}{\sqrt{2} G_F} (1 + \Delta r)$$

- ▶ Precise value sensitive to loop-corrections: QED, top quark, Higgs boson
- ▶ Meanwhile, the indirect SM prediction of the W boson mass has become very precise and a great place to search for the indirect BSM search

$$\begin{aligned} M_W &= 80.3535 \pm 0.0027_{m_t} \pm 0.0030_{\delta_{\text{theo}} m_t} \pm 0.0026_{M_Z} \pm 0.0026_{\alpha_S} \\ &\quad \pm 0.0024_{\Delta \alpha_{\text{had}}} \pm 0.0001_{M_H} \pm 0.0040_{\delta_{\text{theo}} M_W} \text{ GeV}, \\ &= 80.354 \pm 0.007_{\text{tot}} \text{ GeV}, \end{aligned}$$




# Results presented today

- ▶ *Compatibility and combination of world  $W$ -boson mass measurements*, [arXiv:2308.09417](https://arxiv.org/abs/2308.09417), subm. to EPJC
- ▶ *Improved  $W$  boson Mass Measurement using  $\sqrt{s} = 7$  TeV Proton-Proton Collisions with the ATLAS Detector*, [ATLAS-CONF-2023-004](https://atlas.conf.cern.ch/2023/004)
- ▶ *Precise measurements of  $W$  and  $Z$  transverse momentum spectra with the ATLAS detector at  $\sqrt{s} = 5.02$  TeV and 13 TeV*, [ATLAS-CONF-2023-028](https://atlas.conf.cern.ch/2023/028)




# How to measure the $W$ -boson mass

- ▶ Ultimate goal:  $\delta m_W$  better than SM prediction, i.e.  $\lesssim 8 \text{ MeV} = 0.01\%$
- ▶ Single  $W$  bosons produced hadron collisions:  $q\bar{q}' \rightarrow W$ , e.g. TeVatron ( $p\bar{p}$ ) and LHC ( $pp$ )




# How to measure the $W$ -boson mass

- ▶ Ultimate goal:  $\delta m_W$  better than SM prediction, i.e.  $\lesssim 8 \text{ MeV} = 0.01\%$
- ▶ Single  $W$  bosons produced hadron collisions:  $q\bar{q}' \rightarrow W$ , e.g. TeVatron ( $p\bar{p}$ ) and LHC ( $pp$ )




- ▶ Huge event samples available (millions), but unfortunately high-energy hadron collisions a priori not a “precision environment” – a tall mountain to climb:
  - ▶ Leptonic final state in Drell–Yan process  $pp \rightarrow \ell\nu$  eliminates the worst problems from strong interactions, but we now have to compensate for the “missing” information from the neutrino
  - ▶  $Z$  production  $pp \rightarrow \ell\ell$  invaluable to constrain models and calibrate detector
  - ▶ Precision QCD and EW calculations and excellent knowledge of PDFs to compute  $Z$ -to- $W$  differences

# A $W \rightarrow e\nu$ candidate



Run Number: 152409, Event Number: 5966801

Date: 2010-04-05 06:54:50 CEST




$W \rightarrow e\nu$  candidate in  
7 TeV collisions

$p_T(e^+) = 34 \text{ GeV}$

$\eta(e^+) = -0.42$


$E_T^{\text{miss}} = 26 \text{ GeV}$

$M_T = 57 \text{ GeV}$







# The transverse view



- ▶ Collision balanced transversely to the beam (but not longitudinally)
- ▶ Lepton transverse momentum  $p_T^\ell$
- ▶ Remainder of the event: “hadronic recoil”  $\vec{u}_T = p_T^W$
- ▶ neutrino inferred using  $\vec{p}_T^{\text{miss}} = -(\vec{p}_T^\ell + \vec{u}_T)$
- ▶ transverse mass  $m_T = \sqrt{2p_T^\ell p_T^{\text{miss}} (1 - \cos \Delta\phi)}$



# The transverse view



L. Aperio Bella

- ▶ Quantities sensitive to  $m_W$  affected by “physics” and “detector effects”, while we eventually need to understand them to 0.1%
- ▶ Some analyses (e.g. ATLAS) split the sample into many categories with  $W^\pm$ ,  $e/\mu$ , forward/central
- ▶ Obviously all these analyses are blinded w.r.t. the final  $m_W$ , sometimes in several steps




# The transverse view





- ▶ Quantities sensitive to  $m_W$  affected by “physics” and “detector effects”, while we eventually need to understand them to 0.1%
- ▶ Some analyses (e.g. ATLAS) split the sample into many categories with  $W^{\pm}$ ,  $e/\mu$ , forward/central
- ▶ Obviously all these analyses are blinded w.r.t. the final  $m_W$ , sometimes in several steps

# The longitudinal view

- ▶ Longitudinal imbalance in  $W$  production in hadron collisions due to variable and unknown momentum fractions  $x_1, x_2$  of initiating quarks: statistically given by Parton Distribution Functions (PDFs)
- ▶ Flavour matters as well: LHC has more heavy flavour contributions compared to TeVatron



# PDFs in a nutshell



- ▶ Input from Deep Inelastic Lepton-Nucleon scattering and other diverse data
- ▶ Last decades saw improvements in theory, input data and fit methodology: a lot of interesting QCD physics, and an art in itself
- ▶ Fit groups: CT, MMHT/MSHT, NNPDF, in addition ABM+, HERA and HERA+LHC analyses
- ▶ In principle, different groups fit mostly the same data with the same theory and provide uncertainties... good enough for  $m_W$  ?

# The complete “physics model”


- ▶ Typical factorisation of five-dimensional DY cross section

$$\frac{d\sigma}{dp_1 p_2} = \frac{d\sigma}{dm} \frac{d\sigma}{dp_T} \frac{d\sigma}{dy} \left[ 1 + \cos^2 \theta_{CS} + \sum_{i=0}^7 A_i(m, p_T, y) f_i(\theta_{CS}, \phi_{CS}) \right]$$

- ▶  $\frac{d\sigma}{dm}$ : the Breit-Wigner resonance that contains  $m_W$
- ▶  $\frac{d\sigma}{dp_T}$ : transverse momentum, typically constrained by  $Z \rightarrow \ell\ell$  + theory, but can also be measured in  $W$  events
  - ▶ Will come back to this in the final part of the talk
- ▶  $\frac{d\sigma}{dy}$ : rapidity dependence given by PDFs & higher order QCD
- ▶ Angular coefficients  $A_i$  assuming spin-1 boson: higher order QCD with some PDF dependence, can be validated in  $Z \rightarrow \ell\ell$

- ▶ Best measurements from hadron colliders: non-trivial correlations in the physics model, hard to preserve the analyses
- ▶ The LHC-TeVatron EW working group took on the charge:
  - ▶ Improved understanding of QCD and PDF effects (and correlations)
  - ▶ Provide a collaboration endorsed world-average of  $m_W$  measurements


2018 – 2020



# The current status of $m_W$ measurements

- ▶ Best measurements from hadron colliders: non-trivial correlations in the physics model, hard to preserve the analyses
- ▶ The LHC-TeVatron EW working group took on the charge:
  - ▶ Improved understanding of QCD and PDF effects (and correlations)
  - ▶ Provide a collaboration endorsed world-average of  $m_W$  measurements

2021






# The current status of $m_W$ measurements

- ▶ Best measurements from hadron colliders: non-trivial correlations in the physics model, hard to preserve the analyses
- ▶ The LHC-TeVatron EW working group took on the charge:
  - ▶ Improved understanding of QCD and PDF effects (and correlations)
  - ▶ Provide a collaboration endorsed world-average of  $m_W$  measurements — failed

2022




# Inputs and Analysis strategy

- ▶ Challenging measurements typically take multiple years to deliver, using tools and theory modelling available at the time
- ▶ Developed an “update procedure”:



$$m_W^{\text{updated}} = \underbrace{m_W^{\text{ref.}}}_{\text{published}} + \underbrace{\delta m_W^{\text{QCD}}}_{\text{Improved predictions, PDF extrapolation for reference PDF}} + \underbrace{\delta m_W^{\text{PDF}}}_{\text{PDF extrapolation}}$$

- ▶  $\delta m_W^{\text{PDF}}$ : Correct measurements to a new, common PDF baseline
- ▶  $\delta m_W^{\text{QCD}}$ : Correct theory “problems” post-hoc, if beyond the quoted uncertainties
- ▶ Need archeology to understand how  $m_W^{\text{ref}}$  was obtained (papers usually wrong...)




# Starting point and Detector Emulation

- ▶ Original QCD tools and PDFs
  - ▶ CDF: Resbos1 (NLO) with CTEQ6M (NLO), corrected post-hoc to NNPDF3.1
  - ▶ D0: Resbos1 (NNLO) with CTEQ6.6 (NLO)
  - ▶ ATLAS: Powheg+Pythia corrected to NNLO with CT10nnlo
  - ▶ LHCb : Powheg+Pythia with NNLO corrections and PDF average of NNPDF3.1,CT18,MSHT20 (NLO)
- ▶ The original detector-level analysis is usually not accessible (LHCb the exception so far) – instead generate large samples (Powheg NLO and NNLO) and apply fast emulation of detector effects
- ▶ Verified to be good enough to assess shifts from changed theory  $\delta m_W$  at better than 1 MeV



- ▶ Uncovered a wrong modelling of decay angular coefficients in ResBos used for TeVatron analyses: correction of about  $\delta m_W = -10$  MeV
- ▶ In addition: inconsistent  $W$  width assumption (D0), distortion/cutoffs in line shape...
- ▶ CDF (unknowingly?) performed a single correction for PDFs and angular coefficient modelling and eventually needs little correction



Coefficient	$m_T$	$p_T^\ell$	$p_T^\nu$
$A_0$	-6.3	-2.6	-9.1
$A_1$	1.1	1.3	0.3
$A_2$	-0.7	0.4	-3.2
$A_3$	-2.1	-4.1	1.0
$A_4$	-1.4	-3.3	-1.6
$A_0 - A_4$	-9.5	-8.4	-12.5
RESBOS2	$-10.2 \pm 1.1$	$-7.6 \pm 1.2$	$-11.8 \pm 1.4$
Difference	$-0.7 \pm 1.1$	$0.8 \pm 1.2$	$0.7 \pm 1.4$

Table 7: Values of  $\delta m_W^{\text{pol}}$  in MeV associated with reweighting each  $A_i$  coefficient from RESBOS-C to RESBOS2 for the CDF detector, as well as the result of a direct fit to RESBOS2. The result of the direct fit is consistent with that of the reweighting.

- ▶ A fact conveniently ingored in all (!) previous  $m_W$  combinations: measurements should be corrected to the same PDF set before combination
- ▶ Effects can easily be of the same order as the quoted PDF uncertainty


PDF set	D0 $p_T^\ell$	D0 $p_T^{\nu}$	CDF $p_T^\ell$	CDF $p_T^{\nu}$	ATLAS $W^+$	ATLAS $W^-$	LHCb
CTEQ6	-17.0	-17.7	0.0	0.0	-	-	-
CTEQ6.6	0.0	0.0	15.0	17.0	-	-	-
CT10	0.4	-1.3	16.0	16.3	0.0	0.0	-
CT14	-9.7	-10.6	5.8	6.8	-1.2	-5.8	1.1
CT18	-8.2	-9.3	7.2	7.7	12.1	-2.3	-6.0
ABMP16	-19.6	-21.5	-1.4	-2.4	-22.5	-3.1	7.7
MMHT2014	-10.4	-12.7	6.1	5.5	-2.6	9.9	-10.8
MSHT20	-13.7	-15.4	3.6	4.1	-20.9	4.5	-2.0
NNPDF3.1	-1.0	-1.2	14.0	15.1	-14.1	-1.8	6.0
NNPDF4.0	6.7	8.1	20.8	24.1	-22.4	6.9	8.3


Table 3: Values of  $\delta m_W^{\text{PDF}}$  in MeV for each PDF set using the  $p_T^\ell$  (all experiments) or  $p_T^{\nu}$  (CDF and D0) distribution, determined using the WJ-MINNLO calculation.

# PDF uncertainties and correlations

- ▶ Uncertainty perfectly reproduced for ATLAS, while published values for CDF (3.9 MeV) and D0 (11 MeV) established to be underestimates
- ▶ Vast difference in uncertainties using different PDF sets
- ▶ Non-trivial PDF correlation pattern across  $p\bar{p}$ ,  $pp$  and central/forward
- ▶ Choose the CT18 PDF set due to the best compatibility with PDF-sensitive data (not shown) – largest uncertainty on  $m_W$

PDF set	D0	CDF	ATLAS	LHCb
CTEQ6	–	14.1	–	–
CTEQ6.6	<b>15.1</b>	–	–	–
CT10	–	–	<b>9.2</b>	–
CT14	13.8	12.4	11.4	10.8
CT18	14.9	13.4	10.0	12.2
ABMP16	4.5	3.9	4.0	3.0
MMHT2014	8.8	7.7	8.8	8.0
MSHT20	9.4	8.5	7.8	6.8
NNPDF3.1	7.7	<b>6.6</b>	7.4	7.0
NNPDF4.0	8.6	7.7	5.3	4.1






All experiments (4 d.o.f.)					
PDF set	$m_W$	$\sigma_{\text{PDF}}$	$\chi^2$	$p(\chi^2, n)$	
ABMP16	$80392.7 \pm 7.5$	3.2	29	0.0008%	
CT14	$80393.0 \pm 10.9$	7.1	16	0.3%	
CT18	$80394.6 \pm 11.5$	7.7	15	0.5%	
MMHT2014	$80398.0 \pm 9.2$	5.8	17	0.2%	
MSHT20	$80395.1 \pm 9.3$	5.8	16	0.3%	
NNPDF3.1	$80403.0 \pm 8.7$	5.3	23	0.1%	
NNPDF4.0	$80403.1 \pm 8.9$	5.3	28	0.001%	

- ▶ After all corrections applied: combination fails for each PDF set

# Conclusion for the Combination





- ▶ Excluding single experiments gives a clear pattern:
    - ▶ Combinations without CDF work regardless of PDF set
- $$m_W = 80369.2 \pm 13.3 \text{ MeV}$$
- ▶ with 91% probability for the CT18 PDF set
  - ▶ The new CDF measurement is incompatible at  $3.6\sigma$ , even though the PDF uncertainty using CT18 is far larger than the published one
  - ▶ Where next?








- ▶ First  $m_W$  measurement at the LHC was published by ATLAS in late 2016:  $m_W = 80370 \pm 19$  MeV
- ▶ At the time proof “it could be done” and tied for best uncertainty with CDF
- ▶ Reanalysis with a Profile Likelihood fit joined across all categories instead of “classic”  $\chi^2$  fits with offset error propagation

- ▶ Expected a reduction of uncertainty, shift in central value of  $O(16$  MeV) possible




Parameters controlling the correlated uncertainties are shifted and constrained: shift in central value, smaller uncertainty, better Data/Prediction ratios





- ▶ Study of PDF dependence: all results lower than previous result, NNPDF again significantly lower
- ▶ Using CT18 set:  $m_W = 80360 \pm 16$  MeV
  - ▶ 15% better uncertainty than previous publication
  - ▶ One also notices this is closer to the SM and further away from CDF...


PDF-Set	$p_T^\ell$ [MeV]	$m_T$ [MeV]	combined [MeV]
CT10	$80355.6^{+15.8}_{-15.7}$	$80378.1^{+24.4}_{-24.8}$	$80355.8^{+15.7}_{-15.7}$
CT14	$80358.0^{+16.3}_{-16.3}$	$80388.8^{+25.2}_{-25.5}$	$80358.4^{+16.3}_{-16.3}$
CT18	$80360.1^{+16.3}_{-16.3}$	$80382.2^{+25.3}_{-25.3}$	$80360.4^{+16.3}_{-16.3}$
MMHT2014	$80360.3^{+15.9}_{-15.9}$	$80386.2^{+23.9}_{-24.4}$	$80361.0^{+15.9}_{-15.9}$
MSHT20	$80358.9^{+13.0}_{-16.3}$	$80379.4^{+24.6}_{-25.1}$	$80356.3^{+14.6}_{-14.6}$
NNPDF3.1	$80344.7^{+15.6}_{-15.5}$	$80354.3^{+23.6}_{-23.7}$	$80345.0^{+15.5}_{-15.5}$
NNPDF4.0	$80342.2^{+15.3}_{-15.3}$	$80354.3^{+22.3}_{-22.4}$	$80342.9^{+15.3}_{-15.3}$




- ▶ Usually LHC delivers maximum luminosity to ATLAS and CMS: average of 20-70 simultaneous  $pp$  collisions
- ▶ Pileup fills the calorimeters with noise and worsens the “hadronic recoil” measurement
- ▶ However, ATLAS took some special datasets at  $\sqrt{s} = 5$  and 13 TeV: direct measurement of  $p_T^W$  (now), bringing back  $m_T$  into the game for  $m_W$  (future)
- ▶ Also profit from the best luminosity measurements ever at a hadron collider:  $\Delta\mathcal{L} \lesssim 1\%$



# ATLAS low-pileup data





9 additional reconstructed vertices







2 additional reconstructed vertices







- Obviously, we had to calibrate the leptons and the hadronic recoil and determine the backgrounds



- Obviously, we had to calibrate the leptons and the hadronic recoil and determine the backgrounds



- Obviously, we had to calibrate the leptons and the hadronic recoil and determine the backgrounds









# Integrated Results

- ▶ Among the most precise cross  $W$  and  $Z$  cross sections at a hadron collider: 1% luminosity,  $\sim 0.5\%$  other systematics
- ▶ Good agreement with NNLO+NNLL QCD predictions from DYTURBO



PDF set	$W^- \rightarrow l\nu$	$W^+ \rightarrow l\nu$	$Z \rightarrow \ell\ell$
Cross-section at 5.02 TeV [pb]			
CT18	1364	2199	320.9
MSHT20	1351	2185	324.3
NNPDF3.1	1381	2232	329.8
Data	$1384 \pm 16$	$2228 \pm 25$	$333.0 \pm 4.1$
Cross-section at 13 TeV [pb]			
CT18	3410	4462	749.8
MSHT20	3397	4457	766.1
NNPDF3.1	3452	4513	771.4
Data	$3486 \pm 38$	$4571 \pm 49$	$780.3 \pm 10.4$

# Differential Results

- ▶ Differential measurement of boson  $p_T$  distributions challenging to unfold with 5 – 10 GeV wide bins
- ▶ Direct cross-check of dilepton and recoil measurement on  $Z$







# Boson $p_T$ measurement: NNLO+NNLL QCD









- ▶ A  $p_T^W$  measurement in  $\sim 7$  GeV bins at 1–2% accuracy
- ▶ Acceptable agreement with NNLO+NNLL QCD prediction

# Boson $p_T$ measurement: NNLO+NNLL QCD




- ▶ A  $p_T^Z$  measurement in  $\sim 3$  GeV bins at  $< 1\%$  accuracy
- ▶ Acceptable agreement with NNLO+NNLL QCD prediction

# Boson $p_T$ measurement: NNLO+NNLL QCD




# Boson $p_T$ measurement: MC generators with Parton showers




- ▶ Huge variability in parton shower MCs
- ▶ Those carefully tuned to  $\sqrt{s} = 7$  TeV ATLAS data and used for  $m_W - \text{Pythia8 AZ}$  – do a good job, especially at  $\sqrt{s} = 5$  TeV

# Conclusions

- ▶ The  $W$  boson mass is among the key observables to constrain Beyond SM physics
- ▶ The experimental situation is not satisfactory: combination of All-CDF has excellent compatibility and similar precision as CDF alone,  $> 3.6\sigma$  experimental discrepancy
- ▶ Preliminary improved ATLAS  $m_W$  reanalysis pushes the experimental measurement further towards the SM and away from CDF
- ▶ New (preliminary) ATLAS results on  $W$  and  $Z$  transverse momentum spectra at 1 – 2% precision using dedicated low-pileup data open the road towards improvements in modelling for future  $m_W$  analyses and a competitive  $m_W$  measurement using  $m_T$  at LHC
- ▶ PDF knowledge appears yet insufficient to go below  $\Delta m_W \sim 10$  MeV: exploit different environments & new data



L. Aperio Bella











June 26, 2022



In 2012, D0 published a measurement of the W boson mass using  $5.3 \text{ fb}^{-1}$  of Tevatron data (Phys. Rev. Lett. **108**, 151804 (2012)), with a subsequent longer description (Phys. Rev. D **89**, 012005 (2014)). This measurement,  $m_W = 80,375 \pm 23 \text{ MeV}$ , remains the official D0 result.

A study of the remaining approximately  $5 \text{ fb}^{-1}$  of data taken between 2009 and 2011 showed that the deterioration of the detector due to radiation damage effects, combined with the higher pileup owing to the increased instantaneous luminosity, precludes a further precision measurement of the W boson mass.

Correction	$\delta m_W^{\text{QCD}}$ [MeV]					
	$p_T^W$ -constrained			No constraint		
	$p_T^\ell$	$m_T$	$p_T^\nu$	$p_T^\ell$	$m_T$	$p_T^\nu$
Invariant mass	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
Rapidity	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
$A_0$	7.6	10.0	15.8	16.0	12.6	19.5
$A_1$	-2.4	-1.9	-1.8	-1.2	-1.6	-1.4
$A_2$	-3.0	-2.6	2.9	-4.2	-3.0	2.3
$A_3$	2.9	1.6	-0.5	3.5	1.8	-0.2
$A_4$	2.4	-0.1	-0.5	0.1	-0.7	-1.0
$A_0 - A_4$	7.6	7.0	16.0	14.1	9.1	18.9
Total	7.6	7.0	16.0	14.1	9.1	18.9
RESBos2	7.3±1.1	8.4±1.0	16.6±1.2	13.9±1.1	10.3±1.0	19.8±1.2
Non-closure	-0.3±1.1	1.4±1.0	0.6±1.2	-0.2±1.1	1.2±1.0	0.9±1.2

Table 5: Effect of reweighting the angular coefficients in the D0 RESBos1 events to those of RESBos2, as well as a direct fit of RESBos1 to RESBos2. Good closure is observed.









	Stat. Unc.	Muon Unc.	Elec. Unc.	Recoil Unc.	Bckg. Unc.	QCD Unc.	EW Unc.	PDF Unc.	Total Unc.
$W^+$	8.9	6.6	8.2	3.1	5.5	8.4	5.4	14.6	23.4
$W^-$	9.7	7.2	7.8	3.3	6.6	8.3	5.3	13.6	23.4
$W^\pm$	6.8	6.6	6.4	2.9	4.5	8.3	5.5	9.2	18.5

[MeV]



# Lepton Universality

- ▶ In the SM the couplings of the leptons  $e, \mu, \tau$  to  $W$  and  $Z$  bosons are all the same, leading to same branching fractions (ignoring different masses — calculable effect)
- ▶ Very precisely measured at LEP for  $Z \rightarrow ee, \mu\mu, \tau\tau$
- ▶ For  $W$  nowadays strongest constraints from LHC (ATLAS) data







# Precision Observables





- ▶ Muons calibrated using high-statistics  $J/\psi \rightarrow \mu\mu$  sample and transferred to electrons via  $E/p$
- ▶ Measurement of  $Z$ -boson mass:  
 $M_Z = 91\,192.0 \pm 6.4(\text{stat}) \pm 4.0(\text{syst})$  MeV in agreement with LEP
- ▶  $W$  and  $Z$  boson production and decay simulated using RESBOS,  $p_T(Z)$  spectrum tuned to  $Z$  data and validated on  $W$
- ▶ Fit to  $m_T$ ,  $p_T^\ell$  and  $p_T^\nu$  for  $W \rightarrow e\nu$  and  $W \rightarrow \mu\nu$







- ▶ Muons calibrated using high-statistics  $J/\psi \rightarrow \mu\mu$  sample and transferred to electrons via  $E/p$
- ▶ Measurement of  $Z$ -boson mass:  
 $M_Z = 91\,192.0 \pm 6.4(\text{stat}) \pm 4.0(\text{syst})$  MeV in agreement with LEP
- ▶  $W$  and  $Z$  boson production and decay simulated using RESBOS,  $p_T(Z)$  spectrum tuned to  $Z$  data and validated on  $W$
- ▶ Fit to  $m_T$ ,  $p_T^\ell$  and  $p_T^\nu$  for  $W \rightarrow e\nu$  and  $W \rightarrow \mu\nu$



- ▶ Muons calibrated using high-statistics  $J/\psi \rightarrow \mu\mu$  sample and transferred to electrons via  $E/p$
- ▶ Measurement of  $Z$ -boson mass:  
 $M_Z = 91\,192.0 \pm 6.4(\text{stat}) \pm 4.0(\text{syst})$  MeV in agreement with LEP
- ▶  $W$  and  $Z$  boson production and decay simulated using RESBOS,  $p_T(Z)$  spectrum tuned to  $Z$  data and validated on  $W$
- ▶ Fit to  $m_T$ ,  $p_T^\ell$  and  $p_T^\nu$  for  $W \rightarrow e\nu$  and  $W \rightarrow \mu\nu$
- ▶ Measurement of  $W$ -boson mass:  
 $M_W = 80\,433.5 \pm 6.4(\text{stat}) \pm 6.9(\text{syst})$  MeV
  - ▶ Factor 2 better precision than any previous result
  - ▶  $7\sigma$  away from the SM EW fit prediction!



# Colliders

