

The NA62 experiment at CERN: recent results and prospects

Evgueni Goudzovski

(University of Birmingham)

goudzovs@cern.ch

Outline:

- 1) Rare kaon decays in the Standard Model and beyond
- 2) $K^+ \rightarrow \pi^+ \nu \nu$ and related measurements with NA62 Run 1 dataset
- 3) Short-term and long-term plans at CERN
- 4) KOTO experiment at J-PARC: $K_L \rightarrow \pi^0 \nu \nu$ measurement
- 5) Other recent NA62 results
- 6) Summary



Particle Physics Seminar
University of Warwick • 15 October 2020




Rare kaon decays: Standard Model and beyond

Introduction: rare kaon decays

Decay	$\Gamma_{\text{SD}}/\Gamma$	Theory err.*	SM BR $\times 10^{11}$	Exp. BR $\times 10^{11}$
$K_L \rightarrow \mu^+ \mu^-$	10%	30%	79 ± 12 (SD)	684 ± 11
$K_L \rightarrow \pi^0 e^+ e^-$	40%	10%	3.2 ± 1.0	< 28 (@ 90% CL)
$K_L \rightarrow \pi^0 \mu^+ \mu^-$	30%	15%	1.5 ± 0.3	< 38
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	90%	4%	8.4 ± 1.0	< 17.8 (as of 2019)
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	$> 99\%$	2%	3.4 ± 0.6	< 300


*Approx. error on LD-subtracted rate excluding parametric contributions

- ❖ FCNC processes dominated by Z-penguin and box diagrams.
- ❖ SM rates related to V_{CKM} with minimal non-parametric uncertainties.
- ❖ Golden modes $K \rightarrow \pi \nu \bar{\nu}$: uniquely clean theoretically.
- ❖ Decays to charged leptons: affected by larger hadronic uncertainties.



K → πνν in the Standard Model

SM: Z-penguin and box diagrams



“Golden modes”: ultra-rare decays, precise SM predictions.

- ❖ Maximum CKM suppression: $\sim (m_t/m_W)^2 |V_{ts}^* V_{td}|$.
- ❖ No long-distance contributions from amplitudes with intermediate photons.
- ❖ Hadronic matrix element extracted from measured $\text{BR}(K_{e3})$ via isospin rotation.

Mode	Expected BR_{SM}	Experimental status
$K^+ \rightarrow \pi^+ \nu \nu$	$(8.4 \pm 1.0) \times 10^{-11}$	$\text{BR} < 17.8 \times 10^{-11}$ at 90% CL (three NA62 candidates, as of 2019)
$K_L \rightarrow \pi^0 \nu \nu$	$(3.4 \pm 0.6) \times 10^{-11}$	$\text{BR} < 300 \times 10^{-11}$ at 90% CL (KOTO 2015 data)

BR_{SM} : Buras et al., JHEP 1511 (2015) 33; tree-level determination of CKM elements


$K \rightarrow \pi \nu \bar{\nu}$ and the unitarity triangle

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (8.39 \pm 0.30) \times 10^{-11} \cdot \left[\frac{|V_{cb}|}{0.0407} \right]^{2.8} \cdot \left[\frac{\gamma}{73.2^\circ} \right]^{0.74}$$

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = (3.36 \pm 0.05) \times 10^{-11} \cdot$$


$$\cdot \left[\frac{|V_{ub}|}{3.88 \times 10^{-3}} \right]^2 \cdot \left[\frac{|V_{cb}|}{0.0407} \right]^2 \cdot \left[\frac{\sin \gamma}{\sin 73.2^\circ} \right]^2$$

Buras et al., JHEP 1511 (2015) 33




- ❖ Dominant uncertainties: CKM parametric; intrinsic theory uncertainties are **O(1%)**.
- ❖ Work to decrease theory uncertainties [*e.g. Christ et al., PRD 100 (2019) 114506*].
- ❖ Measurements of both K^+ and K_L decays: a clean **$\sin(2\beta)$** measurement, an independent CKM unitarity test.
- ❖ Complementarity to measurements in the **B**-sector. Over-constraining the CKM matrix: reveal the nature of new physics.

CKM unitarity triangle with kaons



$K \rightarrow \pi \nu \bar{\nu}$ and new physics

- ❖ Correlations between BSM contributions K^+ and K_L BRs. [*JHEP 1511 (2015) 166*]
- ❖ Need to measure both K^+ and K_L to discriminate among BSM scenarios.
- ❖ Correlations with other observables (ϵ'/ϵ , ΔM_K , B decays). [*arXiv:2006.01138*]




- ❖ **Green:** models with CKM-like flavour structure
 - ✓ Models with MFV
- ❖ **Blue:** models with new flavour-violating interactions in which LH or RH couplings dominate
 - ✓ **Z'** models with pure LH/RH couplings
- ❖ **Red:** general NP models without the above constraints
- ❖ **The Grossman-Nir bound:** a model-independent relation

$$\frac{\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})} \times \frac{\tau_+}{\tau_L} \leq 1$$

The NA62 experiment at CERN

Kaon programme at CERN




Main **NA62** goal: $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ measurement to **10%** precision with a novel decay-in-flight technique.
 Currently **~300** participants from **31** institutions.

Earlier: NA31

NA48 discovery of direct CPV	1997: $\epsilon'/\epsilon: K_L + K_S$
	1998: $K_L + K_S$
	1999: $K_L + K_S$ K_S HI
	2000: K_L only K_S HI
	2001: $K_L + K_S$ K_S HI
NA48/1	2002: K_S /hyperons
NA48/2	2003: K^+ / K^-
	2004: K^+ / K^-
NA62 R_K run	2007: $K_{e2}^\pm / K_{\mu2}^\pm$ tests
	2008: $K_{e2}^\pm / K_{\mu2}^\pm$ tests
NA62	2015: commissioning
	2016-18: physics run 1
	2021-: physics run 2

Beamline & detector

Un-separated hadron ($p/\pi^+/K^+$) beam.
SPS protons: 400 GeV, nominally $3.3 \times 10^{12}/\text{spill}$.
 K^+ : 75 GeV/c ($\pm 1\%$), divergence $< 100 \mu\text{rad}$.
Nominal beam rate: 750 MHz, K^+ rate: 45 MHz;
 ~ 5 MHz K^+ decays in fiducial volume




- ❖ Currently, 1 year of operation $\approx 10^{18}$ protons on target; 4×10^{12} K^+ decays.
- ❖ Single event sensitivities for K^+ decays: down to $\text{BR} \sim 10^{-12}$.
- ❖ Kinematic rejection factors: 1×10^{-3} for $K^+ \rightarrow \pi^+ \pi^0$, 3×10^{-4} for $K \rightarrow \mu^+ \nu$.
- ❖ Hermetic photon veto: $\pi^0 \rightarrow \gamma \gamma$ decay suppression (for $E_{\pi^0} > 40$ GeV) $\sim 10^{-8}$.
- ❖ Particle ID (RICH+LKr+HAC+MUV): $\sim 10^{-8}$ muon suppression.

NA62 status: Run 1 completed



Run 1 integrated luminosity



2.2×10^{18} POT collected

- ❖ Commissioning run **2015**: minimum bias data ($\sim 3 \times 10^{10}$ protons/pulse).
- ❖ Physics run **2016** (30 days, $\sim 1.3 \times 10^{12}$ ppp): 2×10^{11} useful K^+ decays.
- ❖ Physics run **2017** (160 days, $\sim 1.9 \times 10^{12}$ ppp): 2×10^{12} useful K^+ decays.
- ❖ Physics run **2018** (217 days, $\sim 2.3 \times 10^{12}$ ppp): 4×10^{12} useful K^+ decays.
- ❖ **Run 2** start after the Long Shutdown 2 in **2021** ($\sim 3 \times 10^{12}$ ppp).


$K^+ \rightarrow \pi^+ \nu \nu$ measurement: NA62 Run 1 data set

Analysis of the 2016 data: PLB791 (2019) 156.

Analysis of the 2017 data: arXiv:2007.08218, accepted by JHEP.

*Full Run 1 (2016–18) data set: first presented at ICHEP 2020,
paper in preparation.*

NA62: $K_{\pi\nu\nu}$ signal regions



Main K^+ decay modes (>90% of BR) rejected kinematically.

Resolution on m_{miss}^2 :
 $\sigma = 1.0 \times 10^{-3} \text{ GeV}^4/\text{c}^2$.


Measured kinematical background suppression:

- ✓ $K^+ \rightarrow \pi^+ \pi^0$: 1×10^{-3} ;
- ✓ $K^+ \rightarrow \mu^+ \nu$: 3×10^{-4} .

Further background suppression:

- ✓ PID (calorimeters & Cherenkov detectors):
 μ suppression 10^{-8} ,
 π efficiency = 64%.
- ✓ Hermetic photon veto:
 $\pi^0 \rightarrow \gamma\gamma$ rejection
factor = 1.4×10^{-8} . **11**

Key parameters: timing, PID





Calorimetric PID
(machine learning approach)

RICH PID
(likelihood analysis)

Muon suppression: 10^{-8} ,
pion ID efficiency: 64%

Photon veto:
see later...



Analysis principle



$$N_{\pi\nu\nu}^{exp} \approx N_{\pi\pi} \epsilon_{trigger} \epsilon_{RV} \frac{A_{\pi\nu\nu}}{A_{\pi\pi}} \frac{Br(\pi\nu\nu)}{Br(\pi\pi)} \implies \text{S.E.S.} = \frac{Br(\pi\nu\nu)}{N_{\pi\nu\nu}^{exp}}$$

- $N_{\pi\nu\nu}^{exp}$: expected number of $K_{\pi\nu\nu}$ events
- $Br(\pi\nu\nu)$: Standard Model $K_{\pi\nu\nu}$ branching ratio (central value)
- $N_{\pi\pi}$: $K^+ \rightarrow \pi^+ \pi^0$ events selected from the **control data**, without photon + multiplicity rejection, corrected for pre-scaling
- ϵ_{RV} : “random veto” $K_{\pi\nu\nu}$ efficiency (photon + multiplicity rejection)
- $\epsilon_{trigger}$: trigger efficiency for $K_{\pi\nu\nu}$ events
- $A_{\pi\nu\nu} (A_{\pi\pi})$: acceptances from simulations (**3%** and **8.5%** for 2017 conditions)
- $Br(\pi\pi)$: PDG branching fraction of the $K^+ \rightarrow \pi^+ \pi^0$ decay

Analysis performed in bins of π^+ momentum and instantaneous beam intensity, separately for four data sets.

Single event sensitivity

Beam intensity measured event-by-event from beam tracker (GTK) time sidebands




❖ Integrated over momentum & intensity, $SES_{2018} = (1.11 \pm 0.07) \times 10^{-11}$.
(main uncertainties: trigger, acceptance, random veto)

❖ Expected number of SM events: $N_{\pi\nu\nu} = BR_{SM}/SES = 7.58 \pm 0.40_{\text{sys}} \pm 0.75_{SM} \cdot 14$

$K_{\pi\nu\nu}$ data after selection (2018)


Signal and control regions are blinded




After background evaluation, **control regions are opened first**, and data are compared with background expectations.

“Conventional” backgrounds

Missing mass spectrum of $\pi^+\pi^0$ events (control data)



The largest background from K^+ decays in the vacuum tank:



($K^+ \rightarrow \mu^+\nu$ is treated similarly)

Data events in the $\pi^+\pi^0$ region after the $K_{\pi\nu\nu}$ selection (including π^0 rejection)


$$N_{\text{BKG}} = N(\pi^+\pi^0) f_{\text{kin}}$$

Expected numbers of $K^+ \rightarrow \pi^+\pi^0$ events in signal regions after $K_{\pi\nu\nu}$ selection


Fraction of $\pi^+\pi^0$ events in signal regions measured from control data

Upstream background: type 1

Upstream background event



Upstream background: type 2




Background summary (2018)

Expected SM signal	$7.58(40)_{\text{syst}}(75)_{\text{ext}}$
$K^+ \rightarrow \pi^+ \pi^0(\gamma)$ IB	0.75(4)
$K^+ \rightarrow \mu^+ \nu_\mu(\gamma)$ IB	0.49(5)
$K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$	0.50(11) (from simulations)
$K^+ \rightarrow \pi^+ \pi^- \pi^+$	0.24(8)
$K^+ \rightarrow \pi^+ \gamma \gamma$	< 0.01
$K^+ \rightarrow l^+ \pi^0 \nu_l$	< 0.001
Upstream background	$3.30^{+0.98}_{-0.73}$
Total background	$5.28^{+0.99}_{-0.74}$


- ❖ The largest background is **not due to K^+ decays in the vacuum tank!**
- ❖ Signal acceptance in **2017** reduced by **~40%** to suppress this background.
- ❖ Half of acceptance loss recuperated in **2018** (a new collimator).
- ❖ **Improved the beamline layout** to suppress upstream background in **2021**.
- ❖ Smaller contributions from inelastic interactions are under study.

Opening the box (2018)

Squared missing mass vs pion momentum



Squared missing mass




The 2018 data set:

Candidates observed: **17**
 Expected background: **5.3 ± 1.0**
 Expected SM events: **7.6**

Plus **3** candidates in **2016+17** data.

Result: full Run 1 data set



Categories:

- ❖ Main **2018** data set (**80%**): six pion momentum bins (**15–45 GeV/c**).
- ❖ Second **2018** data sample (old collimator), **2017** and **2016** samples: three separate categories, integrated over pion momentum.


Final result (full Run 1 sample):

$$Br(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (11.0_{-3.5}^{+4.0}{}_{stat.} \pm 0.3_{syst.}) \times 10^{-11}$$

(**3.5 σ** significance)


$K^+ \rightarrow \pi^+ \nu \bar{\nu}$: historical perspective

Time evolution of $BR(K^+ \rightarrow \pi^+ \nu \bar{\nu})$



Hidden-sector physics with $K^+ \rightarrow \pi^+ \nu \nu$


Squared missing mass (2017 data)




- ❖ Signal regions **R1, R2**: search for $K^+ \rightarrow \pi^+ X$ (X =invisible), $0 \leq m_X \leq 100 \text{ MeV}/c^2$ and $160 \leq m_X \leq 260 \text{ MeV}/c^2$.
 - ✓ Interpretation: dark scalar, ALP, QCD axion, axiflavor.
 - ✓ Main background: $K^+ \rightarrow \pi^+ \nu \nu$.
- ❖ The $\pi^+ \pi^0$ region: search for $\pi^0 \rightarrow \text{invisible}$.
 - ✓ SM rate: $\text{BR}(\pi^0 \rightarrow \nu \nu) \sim 10^{-24}$.
 - ✓ Observation = BSM physics.
 - ✓ Reduction of $\pi^0 \rightarrow \gamma \gamma$ background: optimized π^+ momentum range.
 - ✓ Extension: $K^+ \rightarrow \pi^+ X$, with m_X between R1 and R2.

Search for $K^+ \rightarrow \pi^+ X$ (2017 data)

Squared missing mass resolution



UL at 90% CL on $\text{BR}(K^+ \rightarrow \pi^+ X)$ vs m_X





- ❖ Signature: a spike in m_{miss}^2 spectrum of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ candidates.
- ❖ Two candidates in 2017 data, consistent with background: no signal.
- ❖ Upper limits of $\text{BR}(K^+ \rightarrow \pi^+ X)$, depending on X mass and lifetime.
- ❖ Region 2: order of magnitude improvement on BNL E949 [*PRD79 (2009) 092004*]
- ❖ Not limited by background: significant improvements soon.

Search for $\pi^0 \rightarrow$ invisible (2017 data)


- ❖ Rejection of ($K^+ \rightarrow \pi^+ \pi^0 (\gamma)$, $\pi^0 \rightarrow \gamma \gamma$) decays: *CERN-EP-2020-193 (14 Oct 2020)*
simulation based on single-photon efficiency measured with $K^+ \rightarrow \pi^+ \pi^0$ decays.
- ❖ Rejection for $K^+ \rightarrow \pi^+ \nu \nu$ ($15 < p_\pi < 35$ GeV/c): $\epsilon = (1.4 \pm 0.1) \times 10^{-8}$.
- ❖ Rejection for $\pi^0 \rightarrow$ invisible search ($25 < p_\pi < 40$ GeV/c): $\epsilon = (2.8_{-2.1}^{+5.0}) \times 10^{-9}$

Search for $\pi^0 \rightarrow$ invisible: (1/3 of the 2017 data set).

- ❖ $K_{\pi \nu \nu}$ trigger and selection used, but $0.015 < m^2_{\text{miss}} < 0.021$ GeV²/c⁴.
- ❖ Expected $\pi^0 \rightarrow \gamma \gamma$ events: 10_{-8}^{+22} , events observed: 12.



Short-term plans: NA62 Run 2



KOTO result
with 2015 data:
 $BR(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 3.0 \times 10^{-9}$
PRL 122 (2019) 021802

NA62 Run 2 (2021–24):

- ❖ Higher beam intensity.
- ❖ Optimized beamline and new veto detectors to reduce the dominant upstream background.
- ❖ Fourth kaon beam tracker station added.
- ❖ Collect **~100** SM candidates in total.


Expected Run 1+2 sensitivity:
 $\delta BR/BR \approx 10\%$

Long-term plans: $K^+ \rightarrow \pi^+ \nu \nu$ at CERN

- ❖ The $K^+ \rightarrow \pi^+ \nu \nu$ decay in-flight technique is firmly established, and is expected to reach an **O(10%)** measurement by **2024**.
- ❖ A possible next step after LS3 (in **~2027**): a $K^+ \rightarrow \pi^+ \nu \nu$ experiment with **x4** beam intensity (present SPS limit), aiming at **~5%** precision.
 - ✓ Challenge: **O(10ps)** time resolution for key detectors to keep random veto under control, while maintaining other performances.

New pixel beam tracker (GTK):


time resolution: **<50 ps** per plane;
pixel size: **<300×300 μm^2** ;
efficiency: **>99%** per plane (incl. fill factor);
material budget : **0.3–0.5% X_0** ;
beam Intensity: **3 GHz** on **30×60 mm^2** ;
peak intensity: **8.0 MHz/ mm^2** .



A current NA62 GTK station

New STRAW spectrometer:


operation in vacuum;
straw length/diameter: **2.2 m/5 mm**;
trailing time resolution: **~6 ns** per straw;
maximum drift time: **~80 ns**;
layout: **~21000** straws (**4** chambers);
material budget: **1.5% X_0** .



A current NA62 STRAW chamber

Long-term plans: $K_L \rightarrow \pi^0 \nu \nu$ at CERN

- ❖ **KLEVER**: a high-energy experiment (10^{19} pot/year) complementary to KOTO.
- ❖ Photons from K_L decays boosted forward: veto coverage only up to **100 mrad**.
- ❖ Roughly the same vacuum tank layout and fiducial volume as NA62.
- ❖ A possible intermediate NA62/KLEVER step: a $K_L \rightarrow \pi^0 \ell^+ \ell^-$ experiment.



Main detector/veto systems:

UV/AFC	Upstream veto/Active final collimator
LAV1-25	Large-angle vetoes (25 stations)
MEC	Main electromagnetic calorimeter
SAC	Small-angle vetoes
CPV	Charged particle veto
PSD	Pre-shower detector




Target sensitivity:

60 SM $K_L \rightarrow \pi^0 \nu \nu$ events with $S/B \sim 1$
 in 5 years of running;
 $\delta BR(K_L \rightarrow \pi^0 \nu \nu) / BR(K_L \rightarrow \pi^0 \nu \nu) \sim 20\%$.


The KOTO experiment at J-PARC

KOTO at J-PARC: $K_L \rightarrow \pi^0 \nu \nu$

- ❖ Primary beam: **30 GeV** protons;
50 kW = 5.5×10^{13} p/5.2 s (in 2019).
- ❖ Neutral “pencil” beam (at **16°**):
 $\langle p(K_L) \rangle = 2.1$ GeV, with **50%**
in the **(0.7–2.4) GeV** range.
- ❖ Beam composition:
 K_L , neutrons, photons.
- ❖ Fiducial decay region length: **3 m**.
- ❖ CsI calorimeter + hermetic photon veto.



KOTO status



2015 run

- ❖ Reached **40 kW** beam power, 3×10^{19} POT collected.
- ❖ Final 2015 result:
 $BR(K_L \rightarrow \pi^0 \nu \nu) < 3.0 \times 10^{-9}$ at **90%** CL.
PRL 122 (2019) 021802

2016–2018 runs

- ❖ Reached **50 kW** beam power, 4×10^{19} POT collected.
- ❖ Preliminary results reported in 2019/20.

2019 run

- ❖ Analysis in progress.


KOTO: 2016–18 data

Preliminary results (N.Shimizu at ICHEP 2020)

Single-event sensitivity:
 $BR_{SES} = 71 \times 10^{-11}$ ($= 20 \times BR_{SM}$)

Main backgrounds:

source		#BG (90% C.L.)	#BG (68% C.L.)
K+/-	$K^\pm \rightarrow \pi^0 \pi^\pm$	0.09 ± 0.09	0.09 ± 0.09
	$K^\pm \rightarrow \pi^0 e^\pm \nu$	0.90 ± 0.27	0.90 ± 0.27
	$K^\pm \rightarrow \pi^0 \mu^\pm \nu$	< 0.21	< 0.12
Neutron	Upstream π^0	0.001 ± 0.001	0.001 ± 0.001
	Hadron cluster	0.02 ± 0.00	0.02 ± 0.00
	CV-pi0	< 0.10	< 0.05
	CV-eta	0.03 ± 0.01	0.03 ± 0.01
Total	central value	1.05 ± 0.28	1.05 ± 0.28



After a blind analysis, **four candidate events** in the signal region.

- ❖ One event demonstrated to be background (timing in a veto counter).
- ❖ Background estimate (revised): 1.05 ± 0.28 events, mainly from K^\pm decays.
- ❖ The result on $BR(K_L \rightarrow \pi^0 \nu \nu)$ is to be reported soon.

Short-term plans: KOTO step-1

Signal: need **20** times more (flux \times acceptance) to reach SM sensitivity.


- ✓ Increase the beam power (**50**→**100 kW**) gradually by **2024**.
- ✓ 8–16 months of additional running planned in **2020–2024**.

Background: need **~10** times improvement in background rejection to obtain **S/B \approx 1**, assuming SM signal rate.


- ✓ Continuing programme of incremental detector upgrades.

Example:

Dual side readout for CsI calorimeter modules installed at end of 2018 run



Resolve γ/n interaction depth by reading light from front CsI face with a SiPM



Long-term plans: KOTO step-2

To reach $O(100)$ signal events:



- ❖ proton beam power above **100 kW**;
- ❖ new neutral beamline at 5° with $\langle p(K_L) \rangle = 5.2 \text{ GeV}/c$;
- ❖ larger fiducial decay volume;
- ❖ complete rebuild of the detector.

Hadron hall extension required:

- ❖ a joint project with nuclear physics community;
- ❖ on the list of KEK future large-scale projects, with medium priority.


Expected sensitivity:

- ❖ signal acceptance: $5 \times$ KOTO step-1;
- ❖ 60 SM events with $S/B \sim 1$ at **100 kW** beam power ($3 \times 10^7 \text{ s}$).



Recent NA62 results beyond the flagship analysis

$K^+ \rightarrow \pi^+ \mu^+ \mu^-$ measurement (Run 1)



28011 candidates ($\sim 0.05\%$ background):
 ~ 10 times the world sample.

Preliminary result:

$$BR(K^+ \rightarrow \pi^+ \mu^+ \mu^-) = (9.27 \pm 0.07_{\text{stat}} \pm 0.08_{\text{syst}} \pm 0.04_{\text{ext}}) \times 10^{-9}.$$

LU test: $K \rightarrow \pi \mu \mu$ vs $K \rightarrow \pi e e$ FF parameters.

NA62 Preliminary

E865, $K_{\pi ee}$ (1999)
 10300 events – statistical error only

NA48/2, $K_{\pi ee}$ (2009)
 7253 events

NA48/2, $K_{\pi\mu\mu}$ (2011)
 3120 events

NA62, $K_{\pi\mu\mu}$ (2020) – this result
 28011 events

Form factor parameter a_+

NA62 Preliminary

E865, $K_{\pi ee}$ (1999)
 10300 events – statistical error only

NA48/2, $K_{\pi ee}$ (2009)
 7253 events


NA48/2, $K_{\pi\mu\mu}$ (2011)
 3120 events

NA62, $K_{\pi\mu\mu}$ (2020) – this result
 28011 events

Form factor parameter b_+


Search for $K^+ \rightarrow \pi^- \mu^+ \mu^+$ decay (2007)

SM selection: $m(\pi^+ \mu^+ \mu^-)$



Candidates observed: **8357**
 Background: **0.07%**
 $BR(K^+ \rightarrow \pi^+ \mu^+ \mu^-) = (0.962 \pm 0.025) \times 10^{-7}$
 K^+ decays in FV: $(7.94 \pm 0.23) \times 10^{11}$


LNV selection: $m(\pi^- \mu^+ \mu^+)$




Expected background: **0.91 ± 0.41 evt**
 Candidates observed: **1**
 $BR(K^+ \rightarrow \pi^- \mu^+ \mu^+) < 4.2 \times 10^{-11}$ at 90% CL
 [PLB797 (2019) 134794]

Search for $K^+ \rightarrow \pi^- e^+ e^+$ decay (2007)

SM selection: $m(\pi^+ e^+ e^-)$




LNV selection: $m(\pi^- e^+ e^+)$



Candidates observed: **2484**
 $BR(K^+ \rightarrow \pi^+ e^+ e^-) = (3.00 \pm 0.09) \times 10^{-7}$
 K^+ decays in FV: $(2.14 \pm 0.07) \times 10^{11}$

Expected background: 0.16 ± 0.03 evt
 Candidates observed: **0**
 $BR(K^+ \rightarrow \pi^- \mu^+ \mu^+) < 2.2 \times 10^{-10}$ at 90% CL
 [PLB797 (2019) 134794]


Search for $K^+ \rightarrow \pi \mu e$ decays (Run 1)



Bkg. Prediction:
 $N_{SR}^{tot} = 1.06 \pm 0.20$

Observe 0
 events in SR

Preliminary result (ICHEP 2020):
 $BR(K^+ \rightarrow \pi^- \mu^+ e^+) < 4.2 \times 10^{-11}$ at 90% CL





Bkg. Prediction:
 $N_{SR}^{tot} = 0.92 \pm 0.34$

Observe 2
 events in SR

Preliminary result (ICHEP 2020):
 $BR(K^+ \rightarrow \pi^+ \mu^- e^+) < 6.6 \times 10^{-11}$ at 90% CL

HNL production search: data sample

- ❖ Triggers used: $K_{\pi\nu\nu}$ for $K^+ \rightarrow e^+N$; Control/400 for $K^+ \rightarrow \mu^+N$.
- ❖ Numbers of K^+ decays in fiducial volume:
 $N_K = (3.52 \pm 0.02) \times 10^{12}$ in positron case; $N_K = (4.29 \pm 0.02) \times 10^9$ in muon case.
- ❖ Squared missing mass: $m_{\text{miss}}^2 = (P_K - P_\ell)^2$, using STRAW and GTK trackers.
- ❖ HNL production signal: **a spike above continuous missing mass spectrum.**




HNL production search: results

PLB807 (2020) 135599

$|U_{\ell 4}|^2$ limits vs m_{HNL} from production searches

$K^+ \rightarrow \ell^+ N$



- ❖ Full 2016–18 data set for $|U_{e4}|^2$, $\sim 1/3$ of the data set for $|U_{\mu 4}|^2$.
- ❖ Improvement over earlier production searches by up to two orders of magnitude in terms of $|U_{\ell 4}|^2$.
- ❖ For $|U_{e4}|^2$, the BBN-allowed range is excluded up to **340 MeV**.
- ❖ For $|U_{\mu 4}|^2$, sensitivity approaches the E949 one; the search extends to **383 MeV**.

Summary

- ❖ Rare **K** decays: unique new-physics probes up to **O(100 TeV)** mass scale.
- ❖ **NA62 Run 1** in **2016–18**: exposure to **6×10^{12}** **K⁺** decays in flight.
- ❖ Many new results; most importantly, first evidence for the **K⁺→π⁺νν** decay: **20** candidates,

$$Br(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (11.0_{-3.5}^{+4.0}{}_{stat.} \pm 0.3_{syst.}) \times 10^{-11} \quad [preliminary]$$

- ❖ Short-term plans for **K→πνν** decays:
 - ✓ NA62 to reach **O(10%)** precision on **BR(K⁺→π⁺νν)** by **2024** with an established decay in flight technique;
 - ✓ KOTO is making significant progress in background reduction, aiming to reach SM sensitivity to **BR(K_L→π⁰νν)** by **2024**.
- ❖ Next-generation kaon experiments:
 - ✓ High-intensity kaon beam facility at CERN:
O(5%) precision on **BR(K⁺→π⁺νν)** followed by a **K_L** experiment;
 - ✓ KOTO step-2 at J-PARC: plans to measure **BR(K_L→π⁰νν)**;
 - ✓ detector technology: synergies with future collider & flavour experiments.